

	<b>Methylphenidate</b>	<b>Dextroamphetamine</b>	<b>Modafinil</b>	<b>Sodium oxybate</b>
<b>Dose</b>	10-60 mg per day	5-60 mg per day	100-400 mg per day	3-9 g split nightly
<b>Schedule</b>	II	II	IV	III (amended schedule)
<b>Pregnancy category</b>	C	C	C	B
<b>Some interactions</b>	Monoamine oxidase inhibitors (MAOIs)	MAOIs	Oral contraceptives	Sedative hypnotics
<b>Side effects</b>	Nervousness, anxiety, insomnia, tachycardia	Palpitations, tachycardia, psychosis, insomnia, dyskinesia, anorexia, dry mouth, constipation/diarrhea	Headache, nausea, nervousness, anxiety, insomnia	Nausea, vomiting, dizziness
<b>Contraindications</b>	Elderly, seizure disorder, tics, MAOIs use, psychosis, liver disease, history of substance abuse	Cardiovascular disease, hypertension, tics, psychosis, MAOIs use, history of substance abuse	Elderly, heart disease, liver disease, history of substance abuse	Succinic semialdehyde dehydrogenase deficiency, alcohol, sedative hypnotics
<b>30-dose supply, as of July 5, 2006</b>	10 mg tablets; \$18.90	5 mg tablets; \$18.90	100 mg tablets; \$168.50	5 g solution; \$424.44

Figure 1. Some medications for narcolepsy-related excessive daytime sleepiness (EDS).

	<b>Tricyclic antidepressants</b>	<b>Selective serotonin reuptake inhibitors</b>	<b>Sodium oxybate</b>	<b>Nonselective serotonin reuptake inhibitors/non-tricyclic antidepressants</b>
<b>Dose</b>	Protriptyline 2.0-2.5 mg per day Imipramine 25-200 mg per day	Fluoxetine 20-60 mg per day Paroxetine 20-40 mg per day	3-9 g split nightly	Venlafaxine 37.5-225 mg per day Duloxetine 40-60 mg per day
<b>Schedule</b>	Nonscheduled	Nonscheduled	III (amended schedule)	Nonscheduled
<b>Pregnancy category</b>	Protriptyline: C, Imipramine: D	Fluoxetine: C, Paroxetine: D	B	C
<b>Some interactions</b>	Anticholinergics, sympathomimetics	Monoamine oxidase inhibitors (MAOIs)	Sedative hypnotics	MAOIs
<b>Side effects</b>	Anticholinergic effects, QT-segment prolongation	Headaches, nervousness, insomnia, dry mouth, rash	Nausea, dizziness, vomiting	Nausea, vomiting, insomnia, anxiety, tremor
<b>Contraindications</b>	Cardiovascular disease, arrhythmias, elderly, MAOIs use	MAOIs use, suicidality, cardiovascular disease	Succinic semialdehyde dehydrogenase deficiency, alcohol, sedative hypnotics	MAOIs use, hypertension, hepatic/renal dysfunction, suicidality
<b>30-dose supply, as of July 5, 2006</b>	Protriptyline 5 mg tablets; \$51.99 Imipramine 25 mg tablets; \$11.84	Fluoxetine 20 mg tablets; \$15.99 Paroxetine 20 mg tablets; \$32.99	5 g solution; \$424.44	Venlafaxine 37.5 mg tablets; \$56.32 Duloxetine 20 mg tablets; \$101.12

Figure 2. Some medications for narcolepsy-related cataplexy.